

Ondansetron

Adjudication Guideline

Rule Category:
Pharmaceutical

Ref: No:
2024-PH-41

Version Control:
Version No.V3.0

Effective Date:
30/09/2025

Revision Date:
12/09/2026

Approved by:
Daman

Responsible:
Pharmaceutical
Standards& Governance

**Related Adjudication
Guidelines:** N/A

Table of Contents

1.	Abstract	3
1.1	For Members.....	3
1.2	For Medical Professionals.....	3
2.	Scope	3
3.	Adjudication policy.....	4
3.1	Eligibility / Coverage Criteria.....	5
3.2	Requirements for Coverage.....	5
3.3	Non-coverage.....	5
4.	Denial Codes.....	7
5.	Appendices	7
5.1	References	7
5.2	Revision History	8

1. Abstract

1.1 For Members

Ondansetron is indicated for the management of nausea and vomiting associated with chemotherapy, radiotherapy, and surgery. It is particularly useful for patients who are at risk of severe nausea or vomiting following these treatments. This medication must be taken under medical supervision.

1.2 For Medical Professionals

Ondansetron is primarily indicated for the management of nausea and vomiting associated with cancer chemotherapy, radiation therapy, and surgical procedures.

2. Scope

The scope of this adjudication rule highlights the medical indications and coverage requirements of Ondansetron drug for all health insurance plans administered by Daman as per policy terms and conditions.

Ondansetron is a selective 5-HT₃ receptor antagonist used primarily to manage nausea and vomiting caused by cancer chemotherapy, radiation therapy, and surgery. It can be administered orally, rectally, intramuscularly or intravenously.

Mechanism of Action:

Receptor Blocking:

Ondansetron selectively inhibits the 5-HT₃ receptors, found in the gastrointestinal tract and the brain's chemoreceptor trigger zone (CTZ).

Peripheral Effects:


By blocking 5-HT₃ receptors in the gut, ondansetron reduces serotonin-induced nausea and vomiting.

Central Effects:

In the CTZ, the ondansetron prevents nausea and vomiting triggered by stimuli like chemotherapy and radiation.

Tissue Selectivity:

With a high affinity for 5-HT₃ receptors, Ondansetron effectively manages nausea and vomiting while minimizing side effects compared to other antiemetics.



3. Adjudication policy

3.1 Eligibility / Coverage Criteria

Ondansetron is intended for use under the guidance of a healthcare provider if medically justified .

3.2 Therapeutic indications

1. Management of Chemotherapy-Induced Nausea and Vomiting (CINV)
2. Management of Radiotherapy-Induced Nausea and Vomiting (RINV)
3. Management of Postoperative Nausea and Vomiting (PONV)
4. Management of nausea and vomiting associated with gastroenteritis and colitis

The Oral form of Ondansetron is **Not** considered as a first-line treatment

5. Cyclic Vomiting Syndrome in Children:

An idiopathic disorder characterized by recurrent, stereotypical bouts of vomiting with intervening periods of normal or baseline health.

6. Nausea and Vomiting in Pregnancy (NVP)

Ondansetron is recommended as second-line treatment, preferably after the first trimester, if other treatments fail.

- Pregnant Patients before 10 Weeks of Gestation:

Use of ondansetron before 10 weeks of gestation is suspected to cause orofacial malformations. It should only be considered when first-line treatments have failed or if the patient develops complications such as dehydration or poor oral intake that could endanger maternal or fetal health — on a case-by-case basis through appeals.

- Pregnant Patients Between 10 to 20 Weeks of Gestation:


Presenting with nausea and vomiting accompanied by symptoms of hypovolemia, following the clinical step therapy algorithm (*see Image 1*).

- Pregnant Patients Beyond 20 Weeks of Gestation:

Coverage for ondansetron will be considered on a case-by-case basis, as most patients experience resolution of nausea and vomiting after 20 weeks. Therapy beyond this period is typically reserved for rare cases requiring medical justification.

Pregnant patients between 10 to 20 weeks of gestation treatment algorithm:

***Image 1**



3.3 Requirements for Coverage

- Kindly code the ICD-10 and the CPT codes to the highest level of specificity

Red flags:

- Ondansetron is not covered for conditions outside its approved indications, including diagnoses such as "vomiting, unspecified" or "vomiting without accompanying nausea" when reported as standalone diagnosis
- Coverage does not extend to patients without a documented risk of moderate to severe nausea and vomiting.
- Use of ondansetron before 10 weeks of gestation is associated with a small increased risk of oral clefts. It should only be considered when first-line treatments have failed or if complications like dehydration or poor oral intake threaten maternal or fetal health.
- Ondansetron may cause cardiac adverse events, including QT prolongation, arrhythmias, and ischemia. It should be used with extreme caution in patients with existing cardiac conditions or a history of arrhythmia, and avoided if the patient is on other medications known to prolong the QT interval.
- No studies have evaluated the use of orally administered ondansetron for prevention or treatment of postoperative nausea and vomiting (PONV) in children aged ≥ 1 month. In such cases, IV administration is recommended.
- Patients presenting to the emergency department with severe, self-limiting nausea and vomiting due to causes such as viral gastroenteritis, acute mountain sickness, palliative care, or other medical conditions should be treated in emergency settings.
- The Oral form of Ondansetron is considered a second-line treatment for nausea and vomiting associated with gastroenteritis and colitis, except in emergency settings or IP

3.4 Non-coverage

- Not covered for visitor plans

4. Denial Codes

- Kindly apply DOH payment rules and regulations and relevant coding manuals for ICD, Drugs.

Code	Code Description
MNEC-004	Service is not clinically indicated based on good clinical practice, without additional supporting diagnoses/activities
CODE-010	Activity/diagnosis inconsistent with clinician specialty
NCOV-001	Service(s) is (are) not covered
AUTH-001	Prior-Approval is required and was not obtained
MNEC-005	Service/supply may be appropriate, but too frequent
MNEC-006	Alternative drug should be used -Step therapy

5. Appendices

5.1 References

[ZOFRAN \(ondansetron hydrochloride\) Label](#)
[Ondansetron 4 mg film-coated tablets - Summary of Product Characteristics \(SmPC\) - \(emc\)](#)
[National Comprehensive Cancer Network \(NCCN\) Guidelines for Antiemesis, Version 1.2021.](#)
<https://doi.org/10.1213/01.ane.0000295230.55439.f4>
[Radiotherapy-induced nausea and vomiting: Prophylaxis and treatment - UpToDate](#)
<https://www.nice.org.uk/advice/esuom34/ifp/chapter/about-ondansetronRCOG>
<https://www.rcog.org.uk/media/y3fen1x1/gtg69-hyperemesis.pdf>
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9545331/>
<https://www.somanz.org/content/uploads/2020/07/NVP-GUIDELINE-1.2.20-1.pdf>
<https://www.nnuh.nhs.uk/publication/download/hyperemesis-gravidarum-q6-v5/>
<https://obgyn.onlinelibrary.wiley.com/doi/epdf/10.1111/1471-0528.17739>
https://www.uptodate.com/contents/nausea-and-vomiting-of-pregnancy-treatment-and-outcome?search=vomiting%20in%20pregnancy&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9545331/>
https://www.uptodate.com/contents/prevention-of-chemotherapy-induced-nausea-and-vomiting-in-adults?search=prevention%20of%20nausea%20vomiting%20chemo&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
https://www.uptodate.com/contents/acquired-long-qt-syndrome-definitions-pathophysiology-and-causes?search=qt%20prolongation%20risk%20factor&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
https://www.uptodate.com/contents/ondansetron-drug-information?search=ondansetron&source=panel_search_result&selectedTitle=1~150&usage_type=panel&kp_tab=drug_general&display_rank=1#F203733
<https://jamanetwork.com/journals/jama/fullarticle/2718793>
[Ondansetron 4 mg film-coated tablets - Summary of Product Characteristics \(SmPC\) - \(emc\) | 10159](#)

5.2 Revision History

Date	Version No.	Change(s)
02.07.2025	V1	Release of V1.0
22.08.2025	V2	Add step therapy algorithm for Pregnant patients between 10 to 20 weeks
12/09/2026	V3	Guideline review/ No changes

Disclaimer

By accessing these Daman Adjudication Guidelines, you acknowledge that you have read and understood the terms of use set out in the disclaimer below:

The information contained in this Adjudication Guideline is intended to outline the procedures of adjudication of medical claims as applied by the National Health Insurance Company – Daman PJSC (hereinafter "Daman"). The Adjudication Guideline is not intended to be comprehensive, should not be used as treatment guidelines and should only be used for the purpose of reference or guidance for adjudication procedures and shall not be construed as conclusive. Daman in no way interferes with the treatment of patient and will not bear any responsibility for treatment decisions interpreted through Daman Adjudication Guideline. Treatment of patient is and remains at all times the sole responsibility of the treating Healthcare Provider. This Adjudication Guideline does not grant any rights or impose obligations on Daman. The Adjudication Guideline and all of the information it contains are provided "as is" without warranties of any kind, whether express or implied which are hereby expressly disclaimed.

Under no circumstances will Daman be liable to any person or business entity for any direct, indirect, special, incidental, consequential, or other damages arising out of any use of, access to, or inability to use or access to, or reliance on this Adjudication Guideline including but without limitation to, any loss of profits, business interruption, or loss of programs or information, even if Daman has been specifically advised of the possibility of such damages. Daman also disclaims all liability for any material contained in other websites linked to Daman website.

This Adjudication Guideline is subject to the laws, decrees, circulars and regulations of Abu Dhabi and UAE. Any information provided herein is general and is not intended to replace or supersede any laws or regulations related to the Adjudication Guideline as enforced in the UAE issued by any governmental entity or regulatory authority, or any other written document governing the relationship between Daman and its contracting parties.

This Adjudication Guideline is developed by Daman and is the property of Daman and may not be copied, reproduced, distributed or displayed by any third party without Daman's express written consent. This Adjudication Guideline incorporates the Current Procedural Terminology (CPT®), which is a registered trademark of the American Medical Association ("AMA") and the CPT codes and descriptions belong to the AMA. Daman reserves the right to modify, alter, amend or obsolete the Adjudication Guideline at any time by providing one month prior notice.